

A



07/27/99

NEW PATENT APPLICATION  
CONTINUATION-IN-PART  
☒ DIVISIONAL

ASSISTANT COMMISSIONER  
FOR PATENTS  
Box Patent Application  
Washington, D.C. 20231

CERTIFICATE OF MAILING BY "EXPRESS MAIL"  
"EXPRESS MAIL" MAILING LABEL NO EL254461893US  
DATE OF DEPOSIT July 27, 1999 I HEREBY  
CERTIFY THAT THIS PAPER OR FEE IS BEING DEPOSITED WITH THE UNITED  
STATES POSTAL SERVICE "EXPRESS MAIL POST OFFICE TO ADDRESSEE"  
SERVICE UNDER 37 C.F.R. 1.10 ON THE DATE INDICATED ABOVE AND IS  
ADDRESSED TO THE ASSISTANT COMMISSIONER FOR PATENTS, WASHINGTON,  
D.C. 20231  
ANGIE VARGAS  
(TYPED OR PRINTED NAME OF PERSON MAILING PAPER OR FEE)  
Angie  
SIGNATURE OF PERSON MAILING PAPER OR FEE

1c542 U.S. PTO  
09/361655



07/27/99

Sir:

Transmitted herewith for filing is the divisional patent application of:

Inventors: Se-Jin Lee & Aurora F. Esquela

For: **GROWTH DIFFERENTIATION FACTOR-12**

This is a request for filing a \_\_\_ continuation ☒ divisional application under 37  
C.F.R. 1.53(b), of pending prior application Serial No. 08/765,662, filed April 28, 1997, now  
pending.

FULL NAME OF FIRST INVENTOR	LAST NAME: Lee	FIRST NAME: Se-Jin	MIDDLE NAME:
CITIZENSHIP	STATE OR FOREIGN COUNTRY: USA		
POST OFFICE ADDRESS	POST OFFICE ADDRESS: 6711 Chokeberry Road	CITY AND STATE: Baltimore, Maryland	ZIP CODE: 21209
FULL NAME OF SECOND INVENTOR	LAST NAME: Esquela	FIRST NAME: Aurora	MIDDLE NAME: F.
CITIZENSHIP	STATE OR FOREIGN COUNTRY: USA		
POST OFFICE ADDRESS	POST OFFICE ADDRESS: 853 West University Parkway, Apt. 1B	CITY AND STATE: Baltimore, Maryland	ZIP CODE: 92210

In re Application of:  
Lee et al.  
App. No.: Unassigned  
Filed: Herewith  
Page 2

PATENT  
Attorney Docket No.: JHU1550-1

1. X Cancel in this application original claims 1-14 and 23-43 of the prior application before calculating the filing fee. (At least one original independent claim must be retained for filing purposes.)
2. X A preliminary amendment is enclosed.

The filing fee has been calculated as shown below:

For	Number Filed		Number Extra		Rate			Fee	
					Small Entity	Other Entity		Small Entity	Other Entity
Total Claims	8	=	0	X	\$9	\$0	=	\$ .00	\$ 0
Independent Claims	1	=	0	X	\$39	\$0	=	\$ .00	0
Multiple Dependent Claims Presented: <u>  </u> Yes <u>X</u> No					\$130	\$0			0
BASIC FEE					\$380	\$760		\$380.00	\$ 0
TOTAL FEE								\$380.00	

3. X The Assistant Commissioner is hereby authorized to charge a payment of a Total Fee of \$760.00 and any other fees associated with this communication or credit any overpayment to Deposit Account No. 07-1895. A duplicate copy of this Transmittal Sheet is enclosed.  
X Any additional filing fees required under 37 C.F.R. 1.16.  
X Any patent application processing fees under 37 C.F.R. 1.17.
4. X A verified statement claiming small entity status was filed in parent application Serial No. 08/765,662, filed April 28, 1997, and such status is still proper.
5. X The prior application is assigned of record to THE JOHN HOPKINS UNIVERSITY SCHOOL OF MEDICINE
6. X The power of attorney in the prior application is to Lisa A. Haile, Registration No. 38,347
7.    Please transfer the drawings from the prior application to the new application.

In re Application of:  
Lee et al.  
App. No.: Unassigned  
Filed: Herewith  
Page 3

PATENT  
Attorney Docket No.: JHU1550-1

8. X A true copy of the prior application as filed is enclosed, including drawings and the Declaration and Power of Attorney filed in parent application, U.S. Serial No. 08/765,662, filed April 28, 1997.
9. X Copies of the Information Disclosure Statement, PTO Form 1449 and Form 892 from parent application are enclosed.
10. X Copy of Transmittal of Corrected Declaration and Declaration.
11. X Permission To Use Sequence Listing submitted December 19, 1997.

Address all future communications to:

Lisa A. Haile, Ph.D.  
GRAY CARY WARE & FREIDENRICH LLP  
4365 Executive Drive, Suite 1600  
San Diego, California 92121-2189  
Telephone: 858-677-1456  
Facsimile: 858-677-1465

The undersigned states that the enclosed application papers comprise a true copy of the prior application as filed.

Respectfully submitted,

Date: 7/27/99



Lisa A. Haile, Ph.D.  
Attorney for Applicant  
Registration No. 38,374

GRAY CARY WARE & FREIDENRICH LLP  
4365 Executive Drive, Suite 1600  
San Diego, CA 92121-2189

Applicant or Patentee: Lee and Esquela  
 Serial or Patent No.: 08/765,662  
 Filed or Issued: 1/10/97  
 For: GROWTH DIFFERENTIATION FACTOR-12

VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS  
 (37 C.F.R. 1.9(f) and 1.27(c)) - SMALL BUSINESS CONCERN

I hereby declare that I am

- ☐ the owner of the small business concern identified below:  
☒ an official of the small business concern empowered to act on behalf of the concern identified below:

Name of Small Business Concern: John Hopkins University School of Medicine

Address of Small Business Concern: 720 Rutland Avenue, Baltimore, MD 21205

I hereby declare that the above identified small business concern qualifies as a small business concern as defined in 13 C.F.R. 121.12, and reproduced in 37 C.F.R. 1.9(d), for purposes of paying reduced fees to the United States Patent and Trademark Office, in that the number of employees of the concern, including those of its affiliates, does not exceed 500 persons. For purposes of this statement, (1) the number of employees of the business concern is the average over the previous fiscal year of the concern of the persons employed on a full-time, part-time or temporary basis during each of the pay periods of the fiscal year, and (2) concerns are affiliates of each other when either, directly or indirectly, one concern controls or has the power to control the other, or a third party or parties controls or has the power to control both.

I hereby declare that rights under contract or law have been conveyed to and remain with the small business concern identified above with regard to the invention, entitled GROWTH DIFFERENTIATION FACTOR-12 by inventor(s) Se-Jin Lee and Aurora F. Esquela described in

- ☐ the specification filed herewith.  
☒ application serial no. 08/765,662, filed 1/10/97.  
☐ patent no. , issued .

If the rights held by the above identified small business concern are not exclusive, each individual, concern or organization having rights to the invention is listed below and no rights to the invention are held by any person, other than the inventor, who would not qualify as an independent inventor under 37 C.F.R. 1.9(c) if that person made the invention, or by any concern which would not qualify as a small business concern under 37 C.F.R. 1.9(d), or a nonprofit organization under 37 C.F.R. 1.9(e). NOTE: Separate verified statements are required from each named person, concern or organization having rights to the invention averring to their status as small entities. (37 C.F.R. 1.27)

Full Name: John Hopkins University School of Medicine

Address: 720 Rutland Avenue, Baltimore, MD 21205

☐ INDIVIDUAL ☒ SMALL BUSINESS CONCERN ☐ NONPROFIT ORGANIZATION

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 C.F.R. 1.28(b))

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent on which this verified statement is directed.

Name: HOWARD W. CALIFANO

Title: Assistant Dean and Director

Address: 720 Rutland Avenue, Baltimore, MD 21205

Signature: [Signature]

Date: 4-23-97

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

In re Application of: )  
LEE et al ) Group Art Unit: Unknown  
Serial No.: Unknown ) Examiner: Unknown  
Filed: Herewith ) Atty Docket No.: JHU1550-1  
Title: GROWTH DIFFERENTIATION FACTOR-12 )

Assistant Commissioner for Patents  
Washington, D.C. 20231

Sir:

**PRELIMINARY AMENDMENT**

In advance of the first examination in this case, Applicants request amendment of the subject application, as follows:

**IN THE SPECIFICATION**

Page 1, line 2, after "This is a", please insert --divisional of application Serial No. 08/765,662, filed on April 28, 1997, now pending, which is a--.

**IN THE CLAIMS**

Please delete Claims 1-14 and 23-43.

Please amend the remaining claims as follows:


15. (Amended) A method of detecting a cell proliferative disorder comprising contacting [the] an antibody [of claim 12] that specifically binds to growth differentiation factor-12 (GDF-12) polypeptide having an amino acid sequence as set forth in SEQ ID NO:12 with a specimen of a subject suspected of having a GDF-12 associated disorder and detecting binding of the antibody.

16. (Unchanged) The method of claim 15, wherein the cell is a liver cell.
17. (Unchanged) The method of claim 15, wherein the detecting is *in vivo*.
18. (Unchanged) The method of claim 17, wherein the antibody is detectably labeled.
19. (Unchanged) The method of claim 18, wherein the detectable label is selected from the group consisting of a radioisotope, a fluorescent compound, a bioluminescent compound and chemiluminescent compound.
20. (Unchanged) The method of claim 15, wherein the detection is *in vitro*.
21. (Unchanged) The method of claim 20, wherein the antibody is detectably labeled.
22. (Unchanged) The method of claim 21, wherein the label is selected from the group consisting of radioisotope, a fluorescent compound, a bioluminescent compound, a chemiluminescent compound and an enzyme.

#### REMARKS

This Preliminary Amendment is submitted in advance of the first examination of the subject application in order to correct minor typographical errors and clarify certain aspects of the claimed subject matter. No new matter has been entered.

Respectfully submitted,



LISA A. HAILE, Ph.D.  
REG. NO. 38,347

Date:

7/27/99

GRAY CARY WARE & FREIDENRICH LLP  
4365 Executive Drive  
Suite 1600, San Diego, CA 92121-2189

Phone: 858/677-1456 - Fax: 858/677-1465

APPLICATION  
FOR  
UNITED STATES LETTERS PATENT

TITLE: GROWTH DIFFERENTIATION FACTOR-12  
APPLICANT: SE-JIN LEE, AURORA F. ESQUELA

"EXPRESS MAIL" Mailing Label Number EMS4216428US

Date of Deposit 10 JANUARY 1997  
I hereby certify under 37 CFR 1.10 that this correspondence is being deposited with the United States Postal Service as "Express Mail Post Office To Addressee" with sufficient postage on the date indicated above and is addressed to the Commissioner of Patents and Trademarks, Washington, D.C. 20231.

Robert Segool  
Robert Segool

## GROWTH DIFFERENTIATION FACTOR-12

This is a continuation-in-part application of U.S. Serial No. 08/274,215, filed on July 13, 1994.

### BACKGROUND OF THE INVENTION

#### 5      1.      *Field of the Invention*

The invention relates generally to growth factors and specifically to a new member of the transforming growth factor beta (TGF- $\beta$ ) superfamily, which is denoted, growth differentiation factor-12 (GDF-12).

#### 2.      *Description of Related Art*

10      The transforming growth factor  $\beta$  (TGF- $\beta$ ) superfamily encompasses a group of structurally-related proteins which affect a wide range of differentiation processes during embryonic development. The family includes, Mullerian inhibiting substance (MIS), which is required for normal male sex development (Behringer, *et al.*, *Nature*, 345:167, 1990), *Drosophila* decapentaplegic (DPP)  
15      gene product, which is required for dorsal-ventral axis formation and morphogenesis of the imaginal disks (Padgett, *et al.*, *Nature*, 325:81-84, 1987), the *Xenopus* Vg-1 gene product, which localizes to the vegetal pole of eggs ((Weeks, *et al.*, *Cell*, 51:861-867, 1987), the activins (Mason, *et al.*, *Biochem, Biophys. Res. Commun.*, 135:957-964, 1986), which can induce the formation  
20      of mesoderm and anterior structures in *Xenopus* embryos (Thomsen, *et al.*, *Cell*, 63:485, 1990), and the bone morphogenetic proteins (BMPs, osteogenin, OP-1) which can induce *de novo* cartilage and bone formation (Sampath, *et al.*, *J. Biol. Chem.*, 265:13198, 1990). The TGF- $\beta$ s can influence a variety of



-2-

differentiation processes, including adipogenesis, myogenesis, chondrogenesis, hematopoiesis, and epithelial cell differentiation (for review, see Massague, *Cell* 49:437, 1987).

5 The proteins of the TGF- $\beta$  family are initially synthesized as a large precursor protein which subsequently undergoes proteolytic cleavage at a cluster of basic residues approximately 110-140 amino acids from the C-terminus. The C-terminal regions, or mature regions, of the proteins are all structurally related and the different family members can be classified into distinct subgroups based on the extent of their homology. Although the homologies  
10 within particular subgroups range from 70% to 90% amino acid sequence identity, the homologies between subgroups are significantly lower, generally ranging from only 20% to 50%. In each case, the active species appears to be a disulfide-linked dimer of C-terminal fragments. Studies have shown that when the pro-region of a member of the TGF- $\beta$  family is coexpressed with a  
15 mature region of another member of the TGF- $\beta$  family, intracellular dimerization and secretion of biologically active homodimers occur (Gray, A., and Maston, A., *Science*, 247:1328, 1990). Additional studies by Hammonds, et al., (*Molec. Endocrin.* 5:149, 1991) showed that the use of the BMP-2 pro-region combined with the BMP-4 mature region led to dramatically improved  
20 expression of mature BMP-4. For most of the family members that have been studied, the homodimeric species has been found to be biologically active, but for other family members, like the inhibins (Ling, et al., *Nature*, 321:779, 1986) and the TGF- $\beta$ s (Cheifetz, et al., *Cell*, 48:409, 1987), heterodimers have also been detected, and these appear to have different biological properties than  
25 the respective homodimers.

Identification of new factors that are tissue-specific in their expression pattern will provide a greater understanding of that tissue's development and function.

-3-

### SUMMARY OF THE INVENTION

5 The present invention provides a cell growth and differentiation factor, GDF-12, a polynucleotide sequence which encodes the factor, and antibodies which are immunoreactive with the factor. This factor appears to relate to various cell proliferative disorders, especially those involving liver cells.

Thus, in one embodiment, the invention provides a method for detecting a cell proliferative disorder of liver origin and which is associated with GDF-12. In another embodiment, the invention provides a method for treating a cell proliferative disorder by suppressing or enhancing GDF-12 activity.

09361636 072799

-4-

**BRIEF DESCRIPTION OF THE DRAWINGS**

FIGURE 1 shows a Northern blot of RNA prepared from adult tissues probed with a murine GDF-12 probe.

5      FIGURE 2 shows the partial nucleotide and predicted amino acid sequence of human GDF-12.

FIGURE 3 shows the full length nucleotide and predicted amino acid sequence of human GDF-12.

10      FIGURE 4 shows amino acid sequence homologies between human GDF-12 and different members of the TGF- $\beta$  superfamily. Numbers represent amino acid sequence identities between GDF-12 and the indicated family member calculated from the first conserved cysteine to the C-terminus.

09361655 072795

## DETAILED DESCRIPTION OF THE INVENTION

5 The present invention provides a growth and differentiation factor, GDF-12 and a polynucleotide sequence encoding GDF-12. GDF-12 is expressed specifically in liver. In one embodiment, the invention provides a method for detection of a cell proliferative disorder of liver origin which is associated with GDF-12 expression. In another embodiment, the invention provides a method for treating a cell proliferative disorder by using an agent which suppresses or enhances GDF-12 activity.

10 The TGF- $\beta$  superfamily consists of multifunctional polypeptides that control proliferation, differentiation, and other functions in many cell types. Many of the peptides have regulatory, both positive and negative, effects on other peptide growth factors. The structural homology between the GDF-12 protein of this invention and the members of the TGF- $\beta$  family, indicates that GDF-12 is a new member of the family of growth and differentiation factors. Based on 15 the known activities of many of the other members, it can be expected that GDF-12 will also possess biological activities that will make it useful as a diagnostic and therapeutic reagent.

20 In particular, the expression pattern of GDF-12 suggests that GDF-12 possesses activities that will make it useful for the treatment of various diseases involving the liver. For example, when GDF-12 functions to stimulate the growth or differentiation of liver cells, GDF-12 could be used for the treatment of disease states in which the function of the liver is compromised, such as in hepatitis or cirrhosis. Although liver tissue has the capacity to regenerate, GDF-12 could potentially accelerate the normal regenerative process or promote the process in disease states in which the regenerative 25 process is suppressed. Similarly, GDF-12 could be useful for maintaining liver cells or tissue in culture prior to transplantation or for stimulating the growth of liver cells following transplantation; in this regard, because liver cells may be used as a vehicle for delivering genes to liver for gene therapy, GDF-12

could be useful for maintaining or expanding liver cells in culture during or after the introduction of particular genes or for stimulating the growth of these cells following transplantation.

5 Alternatively, when GDF-12 functions as a growth inhibitor, GDF-12 could be used to create cell proliferative disorders involving liver cells, such as hepatocellular carcinoma. Indeed, one member of this superfamily, namely, inhibin alpha, has been shown to function as a tumor suppressor gene, and another member of this superfamily, namely, Mullerian inhibiting substance, has been shown to be capable of inhibiting the growth of tumor cells both *in vitro* and *in vivo*.  
10

This high specificity of GDF-12 expression also suggests potential applications of GDF-12 as a diagnostic tool. In particular, because GDF-12 encodes a secreted factor, levels of GDF-12 could be used to monitor liver function or to detect the presence of neoplasms involving liver cells. In this regard, another  
15 member of this family, namely, inhibin, has been shown to be useful as a marker for ovarian granulosa cell tumors.

The term "substantially pure" as used herein refers to GDF-12 which is substantially free of other proteins, lipids, carbohydrates or other materials with which it is naturally associated. One skilled in the art can purify GDF-12  
20 using standard techniques for protein purification. The substantially pure polypeptide will yield a single major band on a non-reducing polyacrylamide gel. The purity of the GDF-12 polypeptide can also be determined by amino-terminal amino acid sequence analysis. GDF-12 polypeptide includes functional fragments of the polypeptide, as long as the activity of GDF-12  
25 remains. Smaller peptides containing the biological activity of GDF-12 are included in the invention.

The invention provides polynucleotides encoding the GDF-12 protein. These polynucleotides include DNA, cDNA and RNA sequences which encode GDF-

-7-

12. It is understood that all polynucleotides encoding all or a portion of GDF-12 are also included herein, as long as they encode a polypeptide with GDF-12 activity. Such polynucleotides include naturally occurring, synthetic, and intentionally manipulated polynucleotides. For example, GDF-12 polynucleotide may be subjected to site-directed mutagenesis. The polynucleotide sequence for GDF-12 also includes antisense sequences. The polynucleotides of the invention include sequences that are degenerate as a result of the genetic code. There are 20 natural amino acids, most of which are specified by more than one codon. Therefore, all degenerate nucleotide sequences are included in the invention as long as the amino acid sequence of GDF-12 polypeptide encoded by the nucleotide sequence is functionally unchanged.

Specifically disclosed herein is a partial cDNA sequence containing the active portion of the human GDF-12 coding sequence. One of skill in the art could now use this partial sequence to isolate the full length clones. The cDNA clone from which this sequence was obtained is likely to contain the entire coding sequence for GDF-12. The disclosed sequence corresponds to the C-terminal region of the GDF-12 polypeptide. The sequence begins with a putative proteolytic cleavage site, RARRR. Cleavage of the polypeptide at this site would generate an active C-terminal fragment 114 amino acids in length with a predicted molecular weight of 12,500.

The C-terminal region of GDF-12 following the putative proteolytic processing site shows significant homology to the known members of the TGF- $\beta$  superfamily. The GDF-12 sequence contains most of the residues that are highly conserved in other family members (see FIGURE 1). Like the TGF- $\beta$ s and inhibin  $\beta$ s, GDF-12 contains an extra pair of cysteine residues in addition to the 7 cysteines found in virtually all other family members. Among the known family members, GDF-12 is most homologous to Inhibin  $\beta$ B (50% sequence identity) (see FIGURE 4).

-8-

Minor modifications of the recombinant GDF-12 primary amino acid sequence may result in proteins which have substantially equivalent activity as compared to the GDF-12 polypeptide described herein. Such modifications may be deliberate, as by site-directed mutagenesis, or may be spontaneous.

5 All of the polypeptides produced by these modifications are included herein as long as the biological activity of GDF-12 still exists. Further, deletion of one or more amino acids can also result in a modification of the structure of the resultant molecule without significantly altering its biological activity. This can lead to the development of a smaller active molecule which would have broader utility. For example, one can remove amino or carboxy terminal amino acids which are not required for GDF-12 biological activity.

15 The nucleotide sequence encoding the GDF-12 polypeptide of the invention includes the disclosed sequence and conservative variations thereof. The term "conservative variation" as used herein denotes the replacement of an amino acid residue by another, biologically similar residue. Examples of conservative variations include the substitution of one hydrophobic residue such as isoleucine, valine, leucine or methionine for another, or the substitution of one polar residue for another, such as the substitution of arginine for lysine, glutamic for aspartic acid, or glutamine for asparagine, and the like. The term "conservative variation" also includes the use of a substituted amino acid in place of an unsubstituted parent amino acid provided that antibodies raised to the substituted polypeptide also immunoreact with the unsubstituted polypeptide.

The polynucleotide encoding GDF-12 includes the nucleotide sequence in FIGURES 2 and 3 (SEQ ID NO:11 and NO:13, respectively), as well as nucleic acid sequences complementary to that sequence. A complementary sequence may include an antisense nucleotide. When the sequence is RNA, the deoxynucleotides A, G, C, and T of FIGURE 2 and 3 are replaced by ribonucleotides A, G, C, and U, respectively. Also included in the invention are fragments of the above-described nucleic acid sequences that are at least 15 bases in length, which is sufficient to permit the fragment to selectively hybridize to DNA that encodes the protein of FIGURES 2 and 3 (SEQ ID NO:12 and NO:14, respectively) under physiological conditions.

DNA sequences of the invention can be obtained by several methods. For example, the DNA can be isolated using hybridization techniques which are well known in the art. These include, but are not limited to: 1) hybridization of genomic or cDNA libraries with probes to detect homologous nucleotide sequences, 2) polymerase chain reaction (PCR) on genomic DNA or cDNA using primers capable of annealing to the DNA sequence of interest, and 3) antibody screening of expression libraries to detect cloned DNA fragments with shared structural features.

Preferably the GDF-12 polynucleotide of the invention is derived from a mammalian organism, and most preferably from a mouse, rat, or human. Screening procedures which rely on nucleic acid hybridization make it possible to isolate any gene sequence from any organism, provided the appropriate probe is available. Oligonucleotide probes, which correspond to a part of the sequence encoding the protein in question, can be synthesized chemically.

This requires that short, oligopeptide stretches of amino acid sequence must be known. The DNA sequence encoding the protein can be deduced from the genetic code, however, the degeneracy of the code must be taken into account. It is possible to perform a mixed addition reaction when the sequence is degenerate. This includes a heterogeneous mixture of denatured double-stranded DNA. For such screening, hybridization is preferably



-10-

performed on either single-stranded DNA or denatured double-stranded DNA. Hybridization is particularly useful in the detection of cDNA clones derived from sources where an extremely low amount of mRNA sequences relating to the polypeptide of interest are present. In other words, by using stringent hybridization conditions directed to avoid non-specific binding, it is possible, for example, to allow the autoradiographic visualization of a specific cDNA clone by the hybridization of the target DNA to that single probe in the mixture which is its complete complement (Wallace, *et al.*, *Nucl. Acid Res.*, 9:879, 1981).

Therefore, given a partial DNA sequence of the gene of interest, one of skill in the art would be able to prepare probes for isolation of a full length cDNA clone, without undue experimentation (see for example, Ausubel, *et al.*, *Current Protocols in Molecular Biology*, Units 6.3-6.4, Greene Publ., 1994; Maniatis, *et al.*, *Molecular Cloning*, Cold Spring Harbor Laboratories, 1982).

The development of specific DNA sequences encoding GDF-12 can also be obtained by: 1) isolation of double-stranded DNA sequences from the genomic DNA; 2) chemical manufacture of a DNA sequence to provide the necessary codons for the polypeptide of interest; and 3) *in vitro* synthesis of a double-stranded DNA sequence by reverse transcription of mRNA isolated from a eukaryotic donor cell. In the latter case, a double-stranded DNA complement of mRNA is eventually formed which is generally referred to as cDNA.

Of the three above-noted methods for developing specific DNA sequences for use in recombinant procedures, the isolation of genomic DNA isolates is the least common. This is especially true when it is desirable to obtain the microbial expression of mammalian polypeptides due to the presence of introns.

The synthesis of DNA sequences is frequently the method of choice when the entire sequence of amino acid residues of the desired polypeptide product is

-11-

known. When the entire sequence of amino acid residues of the desired polypeptide is not known, the direct synthesis of DNA sequences is not possible and the method of choice is the synthesis of cDNA sequences. Among the standard procedures for isolating cDNA sequences of interest is the formation of plasmid- or phage-carrying cDNA libraries which are derived from reverse transcription of mRNA which is abundant in donor cells that have a high level of genetic expression. When used in combination with polymerase chain reaction technology, even rare expression products can be cloned. In those cases where significant portions of the amino acid sequence of the polypeptide are known, the production of labeled single or double-stranded DNA or RNA probe sequences duplicating a sequence putatively present in the target cDNA may be employed in DNA/DNA hybridization procedures which are carried out on cloned copies of the cDNA which have been denatured into a single-stranded form (Jay, *et al.*, *Nucl. Acid Res.*, 11:2325, 1983).

A cDNA expression library, such as lambda gt11, can be screened indirectly for GDF-12 peptides having at least one epitope, using antibodies specific for GDF-12. Such antibodies can be either polyclonally or monoclonally derived and used to detect expression product indicative of the presence of GDF-12 cDNA.

DNA sequences encoding GDF-12 can be expressed *in vitro* by DNA transfer into a suitable host cell. "Host cells" are cells in which a vector can be propagated and its DNA expressed. The term also includes any progeny of the subject host cell. It is understood that all progeny may not be identical to the parental cell since there may be mutations that occur during replication. However, such progeny are included when the term "host cell" is used. Methods of stable transfer, meaning that the foreign DNA is continuously maintained in the host, are known in the art.

-12-

5 In the present invention, the GDF-12 polynucleotide sequences may be inserted into a recombinant expression vector. The term "recombinant expression vector" refers to a plasmid, virus or other vehicle known in the art that has been manipulated by insertion or incorporation of the GDF-12 genetic sequences. Such expression vectors contain a promoter sequence which facilitates the efficient transcription of the inserted genetic sequence of the host. The expression vector typically contains an origin of replication, a promoter, as well as specific genes which allow phenotypic selection of the transformed cells. Vectors suitable for use in the present invention include, but are not limited to the T7-based expression vector for expression in bacteria (Rosenberg, *et al.*, *Gene*, 56:125, 1987), the pMSXND expression vector for expression in mammalian cells (Lee and Nathans, *J. Biol. Chem.*, 263:3521, 1988) and baculovirus-derived vectors for expression in insect cells. The DNA segment can be present in the vector operably linked to regulatory elements, for example, a promoter (e.g., T7, metallothionein I, or polyhedrin promoters).

Polynucleotide sequences encoding GDF-12 can be expressed in either prokaryotes or eukaryotes. Hosts can include microbial, yeast, insect and mammalian organisms. Methods of expressing DNA sequences having eukaryotic or viral sequences in prokaryotes are well known in the art. Biologically functional viral and plasmid DNA vectors capable of expression and replication in a host are known in the art. Such vectors are used to incorporate DNA sequences of the invention. Preferably, the mature C-terminal region of GDF-12 is expressed from a cDNA clone containing the entire coding sequence of GDF-12. Alternatively, the C-terminal portion of GDF-12 can be expressed as a fusion protein with the pro- region of another member of the TGF- $\beta$  family or co-expressed with another pro- region (see for example, Hammonds, *et al.*, *Molec. Endocrin.* 5:149, 1991; Gray, A., and Mason, A., *Science*, 247:1328, 1990).

-13-

Transformation of a host cell with recombinant DNA may be carried out by conventional techniques as are well known to those skilled in the art. Where the host is prokaryotic, such as *E. coli*, competent cells which are capable of DNA uptake can be prepared from cells harvested after exponential growth phase and subsequently treated by the  $\text{CaCl}_2$  method using procedures well known in the art. Alternatively,  $\text{MgCl}_2$  or  $\text{RbCl}$  can be used. Transformation can also be performed after forming a protoplast of the host cell if desired.

When the host is a eukaryote, such methods of transfection of DNA as calcium phosphate co-precipitates, conventional mechanical procedures such as microinjection, electroporation, insertion of a plasmid encased in liposomes, or virus vectors may be used. Eukaryotic cells can also be cotransformed with DNA sequences encoding the GDF-12 of the invention, and a second foreign DNA molecule encoding a selectable phenotype, such as the herpes simplex thymidine kinase gene. Another method is to use a eukaryotic viral vector, such as simian virus 40 (SV40) or bovine papilloma virus, to transiently infect or transform eukaryotic cells and express the protein. (see for example, *Eukaryotic Viral Vectors*, Cold Spring Harbor Laboratory, Gluzman ed., 1982).

Isolation and purification of microbial expressed polypeptide, or fragments thereof, provided by the invention, may be carried out by conventional means including preparative chromatography and immunological separations involving monoclonal or polyclonal antibodies.

The GDF-12 polypeptides of the invention can also be used to produce antibodies which are immunoreactive or bind to epitopes of the GDF-12 polypeptides. Antibody which consists essentially of pooled monoclonal antibodies with different epitopic specificities, as well as distinct monoclonal antibody preparations are provided. Monoclonal antibodies are made from antigen containing fragments of the protein by methods well known in the art

-14-

(Kohler, *et al.*, *Nature*, 256:495, 1975; *Current Protocols in Molecular Biology*, Ausubel, *et al.*, ed., 1989).

The term "antibody" as used in this invention includes intact molecules as well as fragments thereof, such as Fab, F(ab')<sub>2</sub>, and Fv which are capable of binding the epitopic determinant. These antibody fragments retain some ability to selectively bind with its antigen or receptor and are defined as follows:

- 5 (1) Fab, the fragment which contains a monovalent antigen-binding fragment of an antibody molecule can be produced by digestion of whole antibody with the enzyme papain to yield an intact light chain and a portion of one heavy chain;
- 10 (2) Fab', the fragment of an antibody molecule can be obtained by treating whole antibody with pepsin, followed by reduction, to yield an intact light chain and a portion of the heavy chain; two Fab' fragments are obtained per antibody molecule;
- 15 (3) (Fab')<sub>2</sub>, the fragment of the antibody that can be obtained by treating whole antibody with the enzyme pepsin without subsequent reduction; F(ab')<sub>2</sub> is a dimer of two Fab' fragments held together by two disulfide bonds;
- 20 (4) Fv, defined as a genetically engineered fragment containing the variable region of the light chain and the variable region of the heavy chain expressed as two chains; and

-15-

- (5) Single chain antibody ("SCA"), defined as a genetically engineered molecule containing the variable region of the light chain, the variable region of the heavy chain, linked by a suitable polypeptide linker as a genetically fused single chain molecule.

- 5 Methods of making these fragments are known in the art. (See for example, Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, New York (1988), incorporated herein by reference).

10 As used in this invention, the term "epitope" means any antigenic determinant on an antigen to which the paratope of an antibody binds. Epitopic determinants usually consist of chemically active surface groupings of molecules such as amino acids or sugar side chains and usually have specific three dimensional structural characteristics, as well as specific charge characteristics.

15 Antibodies which bind to the GDF-12 polypeptide of the invention can be prepared using an intact polypeptide or fragments containing small peptides of interest as the immunizing antigen. The polypeptide or a peptide used to immunize an animal can be derived from translated cDNA or chemical synthesis which can be conjugated to a carrier protein, if desired. Such commonly used carriers which are chemically coupled to the peptide include  
20 keyhole limpet hemocyanin (KLH), thyroglobulin, bovine serum albumin (BSA), and tetanus toxoid. The coupled peptide is then used to immunize the animal (e.g., a mouse, a rat, or a rabbit).

25 If desired, polyclonal or monoclonal antibodies can be further purified, for example, by binding to and elution from a matrix to which the polypeptide or a peptide to which the antibodies were raised is bound. Those of skill in the art will know of various techniques common in the immunology arts for purification and/or concentration of polyclonal antibodies, as well as monoclo-

-16-

nal antibodies (See for example, Coligan, *et al.*, Unit 9, *Current Protocols in Immunology*, Wiley Interscience, 1991, incorporated by reference).

It is also possible to use the anti-idiotypic technology to produce monoclonal antibodies which mimic an epitope. For example, an anti-idiotypic monoclonal antibody made to a first monoclonal antibody will have a binding domain in the hypervariable region which is the "image" of the epitope bound by the first monoclonal antibody.

The term "cell-proliferative disorder" denotes malignant as well as non-malignant cell populations which often appear to differ from the surrounding tissue both morphologically and genotypically. Malignant cells (i.e. cancer) develop as a result of a multistep process. The GDF-12 polynucleotide that is an antisense molecule is useful in treating malignancies of the various organ systems, particularly, for example, cells in liver tissue. Essentially, any disorder which is etiologically linked to altered expression of GDF-12 could be considered susceptible to treatment with a GDF-12 suppressing reagent. One such disorder is a malignant cell proliferative disorder, for example.

The invention provides a method for detecting a cell proliferative disorder of muscle or adipose tissue which comprises contacting an anti-GDF-12 antibody with a cell suspected of having a GDF-12 associated disorder and detecting binding to the antibody. The antibody reactive with GDF-12 is labeled with a compound which allows detection of binding to GDF-12. For purposes of the invention, an antibody specific for GDF-12 polypeptide may be used to detect the level of GDF-12 in biological fluids and tissues. Any specimen containing a detectable amount of antigen can be used. A preferred sample of this invention is liver tissue. The level of GDF-12 in the suspect cell can be compared with the level in a normal cell to determine whether the subject has a GDF-12-associated cell proliferative disorder. Preferably the subject is human.

-17-

5 The antibodies of the invention can be used in any subject in which it is desirable to administer *in vitro* or *in vivo* immunodiagnosis or immunotherapy. The antibodies of the invention are suited for use, for example, in immunoassays in which they can be utilized in liquid phase or bound to a solid phase carrier. In addition, the antibodies in these immunoassays can be detectably labeled in various ways. Examples of types of immunoassays which can utilize antibodies of the invention are competitive and non-competitive immunoassays in either a direct or indirect format. Examples of such immunoassays are the radioimmunoassay (RIA) and the sandwich (immunometric) assay. Detection of the antigens using the antibodies of the invention can be done utilizing immunoassays which are run in either the forward, reverse, or simultaneous modes, including immunohistochemical assays on physiological samples. Those of skill in the art will know, or can readily discern, other immunoassay formats without undue experimentation.

10 15 The antibodies of the invention can be bound to many different carriers and used to detect the presence of an antigen comprising the polypeptide of the invention. Examples of well-known carriers include glass, polystyrene, polypropylene, polyethylene, dextran, nylon, amylases, natural and modified celluloses, polyacrylamides, agaroses and magnetite. The nature of the carrier can be either soluble or insoluble for purposes of the invention. Those skilled in the art will know of other suitable carriers for binding antibodies, or will be able to ascertain such, using routine experimentation.

20 25 There are many different labels and methods of labeling known to those of ordinary skill in the art. Examples of the types of labels which can be used in the present invention include enzymes, radioisotopes, fluorescent compounds, colloidal metals, chemiluminescent compounds, phosphorescent compounds, and bioluminescent compounds. Those of ordinary skill in the art will know of other suitable labels for binding to the antibody, or will be able to ascertain such, using routine experimentation.



-18-

Another technique which may also result in greater sensitivity consists of coupling the antibodies to low molecular weight haptens. These haptens can then be specifically detected by means of a second reaction. For example, it is common to use such haptens as biotin, which reacts with avidin, or dinitrophenyl, puridoxal, and fluorescein, which can react with specific anti-hapten antibodies.

In using the monoclonal antibodies of the invention for the *in vivo* detection of antigen, the detectably labeled antibody is given a dose which is diagnostically effective. The term "diagnostically effective" means that the amount of detectably labeled monoclonal antibody is administered in sufficient quantity to enable detection of the site having the antigen comprising a polypeptide of the invention for which the monoclonal antibodies are specific.

The concentration of detectably labeled monoclonal antibody which is administered should be sufficient such that the binding to those cells having the polypeptide is detectable compared to the background. Further, it is desirable that the detectably labeled monoclonal antibody be rapidly cleared from the circulatory system in order to give the best target-to-background signal ratio.

As a rule, the dosage of detectably labeled monoclonal antibody for *in vivo* diagnosis will vary depending on such factors as age, sex, and extent of disease of the individual. Such dosages may vary, for example, depending on whether multiple injections are given, antigenic burden, and other factors known to those of skill in the art.

For *in vivo* diagnostic imaging, the type of detection instrument available is a major factor in selecting a given radioisotope. The radioisotope chosen must have a type of decay which is detectable for a given type of instrument. Still another important factor in selecting a radioisotope for *in vivo* diagnosis is that deleterious radiation with respect to the host is minimized. Ideally, a radio-

-19-

isotope used for *in vivo* imaging will lack a particle emission, but produce a large number of photons in the 140-250 keV range, which may readily be detected by conventional gamma cameras.

For *in vivo* diagnosis radioisotopes may be bound to immunoglobulin either directly or indirectly by using an intermediate functional group. Intermediate functional groups which often are used to bind radioisotopes which exist as metallic ions to immunoglobulins are the bifunctional chelating agents such as diethylenetriaminepentaacetic acid (DTPA) and ethylenediaminetetraacetic acid (EDTA) and similar molecules. Typical examples of metallic ions which can be bound to the monoclonal antibodies of the invention are  $^{111}\text{In}$ ,  $^{97}\text{Ru}$ ,  $^{67}\text{Ga}$ ,  $^{68}\text{Ga}$ ,  $^{72}\text{As}$ ,  $^{89}\text{Zr}$ , and  $^{201}\text{Tl}$ .

The monoclonal antibodies of the invention can also be labeled with a paramagnetic isotope for purposes of *in vivo* diagnosis, as in magnetic resonance imaging (MRI) or electron spin resonance (ESR). In general, any conventional method for visualizing diagnostic imaging can be utilized. Usually gamma and positron emitting radioisotopes are used for camera imaging and paramagnetic isotopes for MRI. Elements which are particularly useful in such techniques include  $^{157}\text{Gd}$ ,  $^{55}\text{Mn}$ ,  $^{162}\text{Dy}$ ,  $^{52}\text{Cr}$ , and  $^{56}\text{Fe}$ .

The monoclonal antibodies of the invention can be used *in vitro* and *in vivo* to monitor the course of amelioration of a GDF-12-associated disease in a subject. Thus, for example, by measuring the increase or decrease in the number of cells expressing antigen comprising a polypeptide of the invention or changes in the concentration of such antigen present in various body fluids, it would be possible to determine whether a particular therapeutic regimen aimed at ameliorating the GDF-12-associated disease is effective. The term "ameliorate" denotes a lessening of the detrimental effect of the GDF-12-associated disease in the subject receiving therapy.

-20-

The present invention identifies a nucleotide sequence that can be expressed in an altered manner as compared to expression in a normal cell, therefore it is possible to design appropriate therapeutic or diagnostic techniques directed to this sequence. Thus, where a cell-proliferative disorder is associated with the expression of GDF-12, nucleic acid sequences that interfere with GDF-12 expression at the translational level can be used. This approach utilizes, for example, antisense nucleic acid and ribozymes to block translation of a specific GDF-12 mRNA, either by masking that mRNA with an antisense nucleic acid or by cleaving it with a ribozyme. Such disorders include liver diseases, for example.

Antisense nucleic acids are DNA or RNA molecules that are complementary to at least a portion of a specific mRNA molecule (Weintraub, *Scientific American*, 262:40, 1990). In the cell, the antisense nucleic acids hybridize to the corresponding mRNA, forming a double-stranded molecule. The antisense nucleic acids interfere with the translation of the mRNA, since the cell will not translate a mRNA that is double-stranded. Antisense oligomers of about 15 nucleotides are preferred, since they are easily synthesized and are less likely to cause problems than larger molecules when introduced into the target GDF-12-producing cell. The use of antisense methods to inhibit the *in vitro* translation of genes is well known in the art (Marcus-Sakura, *Anal.Biochem.*, 172:289, 1988).

Ribozymes are RNA molecules possessing the ability to specifically cleave other single-stranded RNA in a manner analogous to DNA restriction endonucleases. Through the modification of nucleotide sequences which encode these RNAs, it is possible to engineer molecules that recognize specific nucleotide sequences in an RNA molecule and cleave it (Cech, *J.Amer.Med. Assn.*, 260:3030, 1988). A major advantage of this approach is that, because they are sequence-specific, only mRNAs with particular sequences are inactivated.

-21-

There are two basic types of ribozymes namely, *tetrahymena*-type (Hasselhoff, *Nature*, 334:585, 1988) and "hammerhead"-type. *Tetrahymena*-type ribozymes recognize sequences which are four bases in length, while "hammerhead"-type ribozymes recognize base sequences 11-18 bases in length. The longer the recognition sequence, the greater the likelihood that the sequence will occur exclusively in the target mRNA species. Consequently, hammerhead-type ribozymes are preferable to *tetrahymena*-type ribozymes for inactivating a specific mRNA species and 18-based recognition sequences are preferable to shorter recognition sequences.

10 The present invention also provides gene therapy for the treatment of cell proliferative or immunologic disorders which are mediated by GDF-12 protein. Such therapy would achieve its therapeutic effect by introduction of the GDF-12 antisense polynucleotide into cells having the proliferative disorder. Delivery of antisense GDF-12 polynucleotide can be achieved using a  
15 recombinant expression vector such as a chimeric virus or a colloidal dispersion system. Especially preferred for therapeutic delivery of antisense sequences is the use of targeted liposomes.

-22-

Various viral vectors which can be utilized for gene therapy as taught herein include adenovirus, herpes virus, vaccinia, or, preferably, an RNA virus such as a retrovirus. Preferably, the retroviral vector is a derivative of a murine or avian retrovirus. Examples of retroviral vectors in which a single foreign gene can be inserted include, but are not limited to: Moloney murine leukemia virus (MoMuLV), Harvey murine sarcoma virus (HaMuSV), murine mammary tumor virus (MuMTV), and Rous Sarcoma Virus (RSV). A number of additional retroviral vectors can incorporate multiple genes. All of these vectors can transfer or incorporate a gene for a selectable marker so that transduced cells can be identified and generated. By inserting a GDF-12 sequence of interest into the viral vector, along with another gene which encodes the ligand for a receptor on a specific target cell, for example, the vector is now target specific. Retroviral vectors can be made target specific by inserting, for example, a polynucleotide encoding a sugar, a glycolipid, or a protein. Preferred targeting is accomplished by using an antibody to target the retroviral vector. Those of skill in the art will know of, or can readily ascertain without undue experimentation, specific polynucleotide sequences which can be inserted into the retroviral genome to allow target specific delivery of the retroviral vector containing the GDF-12 antisense polynucleotide.

Since recombinant retroviruses are defective, they require assistance in order to produce infectious vector particles. This assistance can be provided, for example, by using helper cell lines that contain plasmids encoding all of the structural genes of the retrovirus under the control of regulatory sequences within the LTR. These plasmids are missing a nucleotide sequence which enables the packaging mechanism to recognize an RNA transcript for encapsidation. Helper cell lines which have deletions of the packaging signal include, but are not limited to  $\Psi$ 2, PA317 and PA12, for example. These cell lines produce empty virions, since no genome is packaged. If a retroviral vector is introduced into such cells in which the packaging signal is intact, but the structural genes are replaced by other genes of interest, the vector can be packaged and vector virion produced.

-23-

Alternatively, NIH 3T3 or other tissue culture cells can be directly transfected with plasmids encoding the retroviral structural genes *gag*, *pol* and *env*, by conventional calcium phosphate transfection. These cells are then transfected with the vector plasmid containing the genes of interest. The  
5 resulting cells release the retroviral vector into the culture medium.

Another targeted delivery system for GDF-12 antisense polynucleotides is a colloidal dispersion system. Colloidal dispersion systems include macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. The  
10 preferred colloidal system of this invention is a liposome. Liposomes are artificial membrane vesicles which are useful as delivery vehicles *in vitro* and *in vivo*. It has been shown that large unilamellar vesicles (LUV), which range in size from 0.2-4.0  $\mu\text{m}$  can encapsulate a substantial percentage of an aqueous buffer containing large macromolecules. RNA, DNA and intact  
15 virions can be encapsulated within the aqueous interior and be delivered to cells in a biologically active form (Fraley, *et al.*, *Trends Biochem. Sci.*, 6:77, 1981). In addition to mammalian cells, liposomes have been used for delivery of polynucleotides in plant, yeast and bacterial cells. In order for a liposome to be an efficient gene transfer vehicle, the following characteristics should be  
20 present: (1) encapsulation of the genes of interest at high efficiency while not compromising their biological activity; (2) preferential and substantial binding to a target cell in comparison to non-target cells; (3) delivery of the aqueous contents of the vesicle to the target cell cytoplasm at high efficiency; and (4) accurate and effective expression of genetic information (Mannino, *et al.*,  
25 *Biotechniques*, 6:682, 1988).

The composition of the liposome is usually a combination of phospholipids, particularly high-phase-transition-temperature phospholipids, usually in combination with steroids, especially cholesterol. Other phospholipids or other lipids may also be used. The physical characteristics of liposomes depend on  
30 pH, ionic strength, and the presence of divalent cations.

-24-

5 Examples of lipids useful in liposome production include phosphatidyl compounds, such as phosphatidylglycerol, phosphatidylcholine, phosphatidylserine, phosphatidylethanolamine, sphingolipids, cerebroside, and gangliosides. Particularly useful are diacylphosphatidylglycerols, where the lipid moiety contains from 14-18 carbon atoms, particularly from 16-18 carbon atoms, and is saturated. Illustrative phospholipids include egg phosphatidylcholine, dipalmitoylphosphatidylcholine and distearoylphosphatidylcholine.

10 The targeting of liposomes can be classified based on anatomical and mechanistic factors. Anatomical classification is based on the level of selectivity, for example, organ-specific, cell-specific, and organelle-specific. Mechanistic targeting can be distinguished based upon whether it is passive or active. Passive targeting utilizes the natural tendency of liposomes to distribute to cells of the reticulo-endothelial system (RES) in organs which contain sinusoidal capillaries. Active targeting, on the other hand, involves alteration of the liposome by coupling the liposome to a specific ligand such as a monoclonal antibody, sugar, glycolipid, or protein, or by changing the composition or size of the liposome in order to achieve targeting to organs and cell types other than the naturally occurring sites of localization.

20 The surface of the targeted delivery system may be modified in a variety of ways. In the case of a liposomal targeted delivery system, lipid groups can be incorporated into the lipid bilayer of the liposome in order to maintain the targeting ligand in stable association with the liposomal bilayer. Various linking groups can be used for joining the lipid chains to the targeting ligand.

25 Due to the expression of GDF-12 in liver tissue, there are a variety of applications using the polypeptide, polynucleotide, and antibodies of the invention, related to these tissues. Such applications include treatment of cell proliferative disorders involving this tissue. In addition, GDF-12 may be useful in various gene therapy procedures.

-25-

The following examples are intended to illustrate but not limit the invention. While they are typical of those that might be used, other procedures known to those skilled in the art may alternatively be used.

### EXAMPLE 1

#### 5 IDENTIFICATION AND ISOLATION OF A NOVEL TGF- $\beta$ FAMILY MEMBER

To identify novel members of the TGF- $\beta$  superfamily, degenerate oligonucleotides were designed which corresponded to two conserved regions among the known family members: one region spanning the two tryptophan residues conserved in most family members and the other region spanning the invariant cysteine residues near the C-terminus. These primers were used for polymerase chain reactions on cDNA synthesized from RNA prepared from whole mouse embryos isolated at day 18.5 of gestation. PCR products were subcloned using restriction sites placed at the 5' ends of the primers, and individual bacterial colonies carrying subcloned inserts were screened by a combination of random sequencing and hybridization analysis to eliminate known members of the superfamily.

GDF-12 was identified from a mixture of PCR products obtained with combinations of primer:

20 SJL218: 5'- CCGGAATTCGGITGG(C/A)G(G/A/T/C)(G/C/A)ATGG  
(A/G)TI(A/G)TITA(T/C)CC (SEQ ID NO:1)

with each of the following 9 primers:

SJL188: 5'- CCGGAATTC(A/G)CAI(C/G)C(A/G)CAIC(C/T)  
(G/A/T/C)(T/A)CIACI(G/A)(T/C)CAT-3' (SEQ ID NO:2)

25 SJL190: 5'- CCGGAATTC(A/G)CAI(C/G)C(A/G)CAIT(C/G)  
(G/A/T/C)(C/T)GIACI(G/A)(T/C)CAT-3' (SEQ ID NO:3)



-26-

- SJL191: 5'- CCGGAATTC(A/G)CAI(C/G)C(A/G)CAIT  
(C/G)(G/AT/C)(T/A)CIACI(G/A)(T/C)CAT-3' (SEQ ID NO:4)
- SJL192: 5'- CCGGAATTC(A/G)CAI(C/G)C(A/G)CAIT(C/G)  
(G/AT/C)(C/G/T)TIACI(G/A)(T/C)CAT-3' (SEQ ID NO:5)
- 5 SJL193: 5'- CCGGAATTC(A/G)CAI(C/G)C(A/G)CAIG  
(A/C)(G/AT/C)(C/T)GIACI(G/A)(T/C)CAT-3' (SEQ ID NO:6)
- SJL194: 5'- CCGGAATTC(A/G)CAI(C/G)C(A/G)CAIG  
(A/C)(G/AT/C)(T/A)CIACI(G/A)(T/C)CAT-3' (SEQ ID NO:7)
- 10 SJL196: 5'- CCGGAATTC(A/G)CAI(C/G)C(A/G)CAI(A/C)G  
(G/AT/C)(C/T)GIACI(G/A)(T/C)CAT-3' (SEQ ID NO:8)
- SJL197: 5'- CCGGAATTC(A/G)CAI(C/G)C(A/G)CAI  
(A/C)G(G/AT/C)(T/A)CIACI(G/A)(T/C)CAT-3' (SEQ ID NO:9)
- SJL198: 5'- CCGGAATTC(A/G)CAI(C/G)C(A/G)CAI(A/C)G  
(G/AT/C)(C/G/T)TIACI(G/A)(T/C)CAT-3' (SEQ ID NO:10)

- 15 PCR using each of these primer combinations was carried out with cDNA prepared from 0.4 $\mu$ g poly A-selected RNA; reactions were carried out at 94°C for 1 minute, 50°C for 2 minutes, and 72°C for 2 minutes for 40 cycles.

- 20 PCR products of approximately 280 base pairs were gel purified, digested with *Eco*RI, gel purified again, and subcloned into the Bluescript vector (Stratagene, San Diego, CA). Bacterial colonies carrying individual subclones were picked into 96 well microtiter plates, and multiple replicas were prepared by plating the cells onto nitrocellulose. The replicate filters were hybridized to probes representing known members of the family, and DNA was prepared from non-hybridizing colonies for sequence analysis.

- 25 Among the colonies analyzed in this manner was one that represented a novel sequence, which was designated GDF-12. This murine sequence was subsequently used to analyze expression patterns and to isolate a human cDNA clone (see below).

-27-

**EXAMPLE 2**  
**EXPRESSION OF GDF-12**

To determine the expression pattern of GDF-12, RNA samples prepared from a variety of adult tissues were screened by Northern analysis. RNA isolation and Northern analysis were carried out as described previously (Lee, S.-J., *Mol. Endocrinol.*, 4:1034, 1990) except that hybridization was carried out in 5X SSPE, 10% dextran sulfate, 50% formamide, 1% SDS, 200  $\mu$ g/ml salmon DNA, and 0.1% each of bovine serum albumin, ficoll, and polyvinylpyrrolidone. Five micrograms of twice poly A-selected RNA prepared from each tissue were electrophoresed on formaldehyde gels, blotted, and probed with GDF-12. As shown in FIGURE 1, the GDF-12 probe detected a single mRNA species approximately 2.8 and 1.9 kb in length, in adult liver.

**EXAMPLE 3**  
**ISOLATION OF cDNA CLONES ENCODING GDF-12**

In order to isolate cDNA clones encoding GDF-11, a cDNA library was prepared in the lambda ZAP II vector (Stratagene) using RNA prepared from human adult liver. From 5  $\mu$ g of twice poly A-selected RNA prepared from human spleen, a cDNA library consisting of 20 million recombinant phage was constructed according to the instructions provided by Stratagene. A portion of this library was screened without amplification using the murine GDF-12 PCR product as a probe. Library screening and characterization of cDNA inserts were carried out as described previously (Lee, *Mol. Endocrinol.*, 4:1034, 1990), except that the final wash was carried out in 2xSSC.

-28-

Partial sequence analysis of the first isolated clone showed that it contained the entire coding sequence of GDF-12. A portion of the nucleotide and predicted amino acid sequence of this clone is shown in FIGURE 2 and SEQ ID NOs: 11 and 12. The sequence begins with a putative proteolytic cleavage site which is followed by a C-terminal region of 114 amino acids. The active C-terminal fragment is predicted to have a molecular weight of approximately 12,500.

The entire nucleotide sequence of the longest human GDF-12 cDNA clone isolated is shown in FIGURE 3 and SEQ ID NO:13. The 2419 base pair sequence contains a single long open reading frame beginning with a methionine codon at nucleotides 218-220 and extending for 350 codons. The sequence contains an in-frame stop codon upstream of the putative initiating methionine. The predicted amino acid sequence (SEQ ID NO:14) contains a stretch of hydrophobic amino acids near the N-terminus suggestive of a signal sequence for secretion, one potential N-linked glycosylation site at amino acids 232-236 (box). The C-terminal region following the putative processing site (shaded box) contains all of the hallmarks present in other TGF- $\beta$  family members (see above).

The C-terminal region following the predicted cleavage site contains all the hallmarks present in other TGF- $\beta$  family members. GDF-12 contains most of the residues that are highly conserved in other family members, including the seven cysteine residues with their characteristic spacing. Like the TGF- $\beta$ 's, and the inhibin  $\beta$ 's, GDF-12 also contains two additional cysteine residues. In the case of TGF- $\beta$ 2, these additional cysteine residues are known to form an intramolecular disulfide bond (Daopin, *et al.*, *Science*, 257:369, 1992; Schlunegger and Grutter, *Nature*, 358:430, 1992). A tabulation of the amino acid sequence homologies between GDF-12 and the other TGF- $\beta$  family members is shown in FIGURE 4. Numbers represent percent amino acid identities between each pair calculated from the first conserved cysteine to the C-terminus.

[illegible]

**THE**  
JOURNAL  
**OF THE**  
ROYAL ANTHROPOLOGICAL INSTITUTE  
AND  
THE  
ETHNOGRAPHIC SOCIETY OF LONDON  
PUBLISHED BY ORDER OF THE SOCIETIES.  
LONDON:  
PRINTED BY WILKINSON AND SONS,  
STATIONERS' HALL COURT, LANCING LANE.

-30-

## SEQUENCE LISTING

## (1) GENERAL INFORMATION

(i) APPLICANT: THE JOHNS HOPKINS UNIVERSITY SCHOOL OF MEDICINE

(ii) TITLE OF THE INVENTION: GROWTH DIFFERENTIATION FACTOR-12

5 (iii) NUMBER OF SEQUENCES: 14

## (iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Fish &amp; Richardson

(B) STREET: 4225 Executive Square, Suite 1400

(C) CITY: La Jolla

10 (D) STATE: CA

(E) COUNTRY: USA

(F) ZIP: 92037

## (v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Diskette

15 (B) COMPUTER: IBM Compatible

(C) OPERATING SYSTEM: DOS

(D) SOFTWARE: FastSEQ Version 1.5

## (vi) CURRENT APPLICATION DATA:

20 (A) APPLICATION NUMBER: PCT/US95/

(B) FILING DATE: 12-JUL-1995

(C) CLASSIFICATION:

## (vii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Haile, Ph.D., Lisa A

(B) REGISTRATION NUMBER: 38,347

25 (C) REFERENCE/DOCKET NUMBER: 07265/042WO1 (FD-3830)

## (ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 619-678-5070

(B) TELEFAX: 619-678-5099

(C) TELEX:

30 (2) INFORMATION FOR SEQ ID NO:1:

-31-

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 34 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(1x) FEATURE:

(A) NAME/KEY: Modified Base

(B) LOCATION: 12...12

(D) OTHER INFORMATION: Inosine

(A) NAME/KEY: Modified Base

(B) LOCATION: 26...26

(D) OTHER INFORMATION: Inosine

(A) NAME/KEY: Modified Base

(B) LOCATION: 29...29

(D) OTHER INFORMATION: Inosine

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CCGGAATTCG GNTGGMGNVA TGGRTNRTNT AYCC

(2) INFORMATION FOR SEQ ID NO:2:

## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(11) MOLECULE TYPE: cDNA

(111) HYPOTHETICAL: NO

(1v) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(1x) FEATURE:

(A) NAME/KEY: Modified Base

-32-

(B) LOCATION: 13...13  
(D) OTHER INFORMATION: Inosine

5

(A) NAME/KEY:Modified Base  
(B) LOCATION: 19...19  
(D) OTHER INFORMATION: Inosine

10

(A) NAME/KEY:Modified Base  
(B) LOCATION: 25...25  
(D) OTHER INFORMATION: Inosine; Inosine also at position  
28

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

CCGGAATTCTCANSRCANCA YNWCNACNRY CAT

33

(2) INFORMATION FOR SEQ ID NO:3:

15

(1) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 33 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: cDNA  
(iii) HYPOTHETICAL: NO  
(iv) ANTISENSE: NO  
(v) FRAGMENT TYPE:  
(vi) ORIGINAL SOURCE:

25

(ix) FEATURE:

(A) NAME/KEY: Modified Base  
(B) LOCATION: 13...13  
(D) OTHER INFORMATION: Inosine

30

(A) NAME/KEY:Modified Base  
(B) LOCATION: 19...19  
(D) OTHER INFORMATION: Inosine

35

(A) NAME/KEY:Modified Base  
(B) LOCATION: 25...25  
(D) OTHER INFORMATION: Inosine; Inosine also at position  
28

-33-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CCGGAATTCTCANSRCANTSNYGNACNRYCAT

33

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

5

- (A) LENGTH: 33 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

10

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(1x) FEATURE:

15

- (A) NAME/KEY: Modified Base
- (B) LOCATION: 13...13
- (D) OTHER INFORMATION: Inosine

20

- (A) NAME/KEY: Modified Base
- (B) LOCATION: 19...19
- (D) OTHER INFORMATION: Inosine

25

- (A) NAME/KEY: Modified Base
- (B) LOCATION: 25...25
- (D) OTHER INFORMATION: Inosine; Inosine also at position

28



-34-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CCGGAATTCT CANSRCANT SNWCNACNRY CAT

33

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

5

- (A) LENGTH: 33 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

10

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(ix) FEATURE:

15

(A) NAME/KEY: Modified Base

(B) LOCATION: 13...13

(D) OTHER INFORMATION: Inosine

20

(A) NAME/KEY: Modified Base

(B) LOCATION: 19...19

(D) OTHER INFORMATION: Inosine

25

(A) NAME/KEY: Modified Base

(B) LOCATION: 25...25

(D) OTHER INFORMATION: Inosine; Inosine also at position

28

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CCGGAATTCT CANSRCANT SNBTNACNRY CAT

33

(2) INFORMATION FOR SEQ ID NO:6:

30

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

-35-

- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE:
- (vi) ORIGINAL SOURCE:
  - (ix) FEATURE:

5 (A) NAME/KEY: Modified Base  
(B) LOCATION: 13...13  
(D) OTHER INFORMATION: Inosine

10 (A) NAME/KEY: Modified Base  
(B) LOCATION: 19...19  
(D) OTHER INFORMATION: Inosine

15 (A) NAME/KEY: Modified Base  
(B) LOCATION: 25...25  
(D) OTHER INFORMATION: Inosine; Inosine also at position  
28

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:6:

CCGGAATTCTCANSRCRCANGMNYGNACNRYCAT

33

(2) INFORMATION FOR SEQ ID NO:7:

20 (1) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 33 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: cDNA  
(iii) HYPOTHETICAL: NO  
(iv) ANTISENSE: NO  
(v) FRAGMENT TYPE:  
(vi) ORIGINAL SOURCE:

30 (ix) FEATURE:  
  
(A) NAME/KEY: Modified Base  
(B) LOCATION: 13...13  
(D) OTHER INFORMATION: Inosine

35 (A) NAME/KEY: Modified Base  
(B) LOCATION: 19...19  
(D) OTHER INFORMATION: Inosine

-36-

- (A) NAME/KEY:Modified Base  
(B) LOCATION: 25...25  
(D) OTHER INFORMATION: Inosine; Inosine also at position  
28

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CCGGAATTCTCANSRCRCANGMNWCNACNRYCAT

33

(2) INFORMATION FOR SEQ ID NO:8:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(1x) FEATURE:

- (A) NAME/KEY: Modified Base  
(B) LOCATION: 13...13  
(D) OTHER INFORMATION: Inosine

- (A) NAME/KEY:Modified Base  
(B) LOCATION: 19...19  
(D) OTHER INFORMATION: Inosine

- (A) NAME/KEY:Modified Base  
(B) LOCATION: 25...25  
(D) OTHER INFORMATION: Inosine; Inosine also at position  
28

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:8:

CCGGAATTCTCANSRCRCANMGNYGACNRYCAT

33

(2) INFORMATION FOR SEQ ID NO:9:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 base pairs  
(B) TYPE: nucleic acid

-37-

(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA  
(iii) HYPOTHETICAL: NO  
(iv) ANTISENSE: NO  
(v) FRAGMENT TYPE:  
(vi) ORIGINAL SOURCE:  
(ix) FEATURE:

(A) NAME/KEY: Modified Base  
(B) LOCATION: 13...13  
(D) OTHER INFORMATION: Inosine

(A) NAME/KEY: Modified Base  
(B) LOCATION: 19...19  
(D) OTHER INFORMATION: Inosine

(A) NAME/KEY: Modified Base  
(B) LOCATION: 25...25  
(D) OTHER INFORMATION: Inosine; Inosine also at position  
28

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

CCGGAATTCTC CAnSCRCANM GNWCNACNMY CAT

33

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 33 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA  
(iii) HYPOTHETICAL: NO  
(iv) ANTISENSE: NO  
(v) FRAGMENT TYPE:  
(vi) ORIGINAL SOURCE:  
(ix) FEATURE:

(A) NAME/KEY: Modified Base  
(B) LOCATION: 13...13  
(D) OTHER INFORMATION: Inosine

-38-

(A) NAME/KEY: Modified Base  
 (B) LOCATION: 19...19  
 (D) OTHER INFORMATION: Inosine

5

(A) NAME/KEY: Modified Base  
 (B) LOCATION: 25...25  
 (D) OTHER INFORMATION: Inosine; Inosine also at position  
 28

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

10

CCGGAATTCTC CANSRCANM GNWCNACNMY CAT

33

(2) INFORMATION FOR SEQ ID NO:11:

(1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 360 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

20

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(ix) FEATURE:

(A) NAME/KEY: Coding Sequence

(B) LOCATION: 1...357

25

(D) OTHER INFORMATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

CGG GCC AGG AGG AGG ACC CCC ACC TGT GAG CCT GCG ACC CCC TTA TGT  
 Arg Ala Arg Arg Arg Thr Pro Thr Cys Glu Pro Ala Thr Pro Leu Cys  
 1 5 10 15

48

30

TGC AGG CGA GAC CAT TAC GTA GAC TTC CAG GAA CTG GGA TGG CGG GAC  
 Cys Arg Arg Asp His Tyr Val Asp Phe Gln Glu Leu Gly Trp Arg Asp  
 20 25 30

96

35

TGG ATA CTG CAG CCC GAG GGG TAC CAG CTG AAT TAC TGC AGT GGG CAG  
 Trp Ile Leu Gln Pro Glu Gly Tyr Gln Leu Asn Tyr Cys Ser Gly Gln  
 35 40 45

144

-39-

TGC CCT CCC CAC CTG GCT GGC AGC CCA GGC ATT GCT GCC TCT TTC CAT 192  
 Cys Pro Pro His Leu Ala Gly Ser Pro Gly Ile Ala Ala Ser Phe His  
 50 55 60

5 TCT GCC GTC TTC AGC CTC CTC AAA GCC AAC AAT CCT TGG CCT GCC AGT 240  
 Ser Ala Val Phe Ser Leu Leu Lys Ala Asn Asn Pro Trp Pro Ala Ser  
 65 70 75 80

ACC TCC TGT TGT GTC CCT ACT GCC CGA AGG CCC CTC TCT CTC CTC TAC 288  
 Thr Ser Cys Cys Val Pro Thr Ala Arg Arg Pro Leu Ser Leu Leu Tyr  
 85 90 95

10 CTG GAT CAT AAT GGC AAT GTG GTC AAG ACG GAT GTG CCA GAT ATG GTG 336  
 Leu Asp His Asn Gly Asn Val Val Lys Thr Asp Val Pro Asp Met Val  
 100 105 110

15 GTG GAG GCC TGT GGC TGC AGC TAG 360  
 Val Glu Ala Cys Gly Cys Ser  
 115

## (2) INFORMATION FOR SEQ ID NO:12:

## (1) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 119 amino acids  
 (B) TYPE: amino acids  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(11) MOLECULE TYPE: protein

(111) HYPOTHETICAL: NO

(1v) ANTISENSE: NO

25 (v) FRAGMENT TYPE: internal

(v1) ORIGINAL SOURCE:

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Arg Ala Arg Arg Arg Thr Pro Thr Cys Glu Pro Ala Thr Pro Leu Cys  
 1 5 10 15

30 Cys Arg Arg Asp His Tyr Val Asp Phe Gln Glu Leu Gly Trp Arg Asp  
 20 25 30

Trp Ile Leu Gln Pro Glu Gly Tyr Gln Leu Asn Tyr Cys Ser Gly Gln  
 35 40 45

35 Cys Pro Pro His Leu Ala Gly Ser Pro Gly Ile Ala Ala Ser Phe His  
 50 55 60

-40-

Ser Ala Val Phe Ser Leu Leu Lys Ala Asn Asn Pro Trp Pro Ala Ser  
65 70 75 80

Thr Ser Cys Cys Val Pro Thr Ala Arg Arg Pro Leu Ser Leu Leu Tyr  
85 90 95

5 Leu Asp His Asn Gly Asn Val Val Lys Thr Asp Val Pro Asp Met Val  
100 105 110

Val Glu Ala Cys Gly Cys Ser  
115

(2) INFORMATION FOR SEQ ID NO:13:

10 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 2419 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: cDNA  
(iii) HYPOTHETICAL: NO  
(iv) ANTISENSE: NO  
(v) FRAGMENT TYPE:  
(vi) ORIGINAL SOURCE:

20 (ix) FEATURE:  
  
(A) NAME/KEY: Coding Sequence  
(B) LOCATION: 218...1267  
(D) OTHER INFORMATION:

25 (x1) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GAGCTGTGAG GGTCAAGCAC AGCTATCCAT CAGATGATCT ACTTTCAGCC TTCCTGAGTC 60

CCAGACAATA GAAGACAGGT GGCTGTACCC TTGGCCAAGG GTAGGTGTGG CAGTGGTGTC 120

TGCTGTCACT GTGCCCTCAT TGGCCCCCAG CAATCAGACT CAACAGACGG AGCAACTGCC 180

ATCCGAGGCT CCTGAACCAG GGCCATTCAC CAGGAGC ATG CGG CTC CCT GAT GTC 235

30 Met Arg Leu Pro Asp Val  
1 5

CAG CTC TGG CTG GTG CTG CTG TGG GCA CTG GTG CGA GCA CAG GGG ACA 283

Gln Leu Trp Leu Val Leu Leu Trp Ala Leu Val Arg Ala Gln Gly Thr  
10 15 20

-41-

	GGG TCT GTG TGT CCC TCC TGT GGG GGC TCC AAA CTG GCA CCC CAA GCA	331
	Gly Ser Val Cys Pro Ser Cys Gly Gly Ser Lys Leu Ala Pro Gln Ala	
	25 30 35	
5	GAA CGA GCT CTG GTG CTG GAG CTA GCC AAG CAG CAA ATC CTG GAT GGG	379
	Glu Arg Ala Leu Val Leu Glu Leu Ala Lys Gln Gln Ile Leu Asp Gly	
	40 45 50	
	TTG CAC CTG ACC AGT CGT CCC AGA ATA ACT CAT CCT CCA CCC CAG GCA	427
	Leu His Leu Thr Ser Arg Pro Arg Ile Thr His Pro Pro Pro Gln Ala	
	55 60 65 70	
10	GCG CTG ACC AGA GCC CTC CGG AGA CTA CAG CCA GGG AGT GTG GCT CCA	475
	Ala Leu Thr Arg Ala Leu Arg Arg Leu Gln Pro Gly Ser Val Ala Pro	
	75 80 85	
	GGG AAT GGG GAG GAG GTC ATC AGC TTT GCT ACT GTC ACA GAC TCC ACT	523
	Gly Asn Gly Glu Glu Val Ile Ser Phe Ala Thr Val Thr Asp Ser Thr	
15	90 95 100	
	TCA GCC TAC AGC TCC CTG CTC ACT TTT CAC CTG TCC ACT CCT CGG TCC	571
	Ser Ala Tyr Ser Ser Leu Leu Thr Phe His Leu Ser Thr Pro Arg Ser	
	105 110 115	
20	CAC CAC CTG TAC CAT GCC CGC CTG TGG CTG CAC GTG CTC CCC ACC CTT	619
	His His Leu Tyr His Ala Arg Leu Trp Leu His Val Leu Pro Thr Leu	
	120 125 130	
	CCT GGC ACT CTT TGC TTG AGG ATC TTC CGA TGG GGA CCA AGG AGG AGG	667
	Pro Gly Thr Leu Cys Leu Arg Ile Phe Arg Trp Gly Pro Arg Arg Arg	
	135 140 145 150	
25	CGC CAA GGG TCC CGC ACT CTC CTG GCT GAG CAC CAC ATC ACC AAC CTG	715
	Arg Gln Gly Ser Arg Thr Leu Leu Ala Glu His His Ile Thr Asn Leu	
	155 160 165	
	GGC TGG CAT ACC TTA ACT CTG CCC TCT AGT GGC TTG AGG GGT GAG AAG	763
	Gly Trp His Thr Leu Thr Leu Pro Ser Ser Gly Leu Arg Gly Glu Lys	
30	170 175 180	
	TCT GGT GTC CTG AAA CTG CAA CTA GAC TGC AGA CCC CTA GAA GGC AAC	811
	Ser Gly Val Leu Lys Leu Gln Leu Asp Cys Arg Pro Leu Glu Gly Asn	
	185 190 195	
35	AGC ACA GTT ACT GGA CAA CCG AGG CGG CTC TTG GAC ACA GCA GGA CAC	859
	Ser Thr Val Thr Gly Gln Pro Arg Arg Leu Leu Asp Thr Ala Gly His	
	200 205 210	

09361633-0712793



	CAG CAG CCC TTC CTA GAG CTT AAG ATC CGA GCC AAT GAG CCT GGA GCA	907
	Gln Gln Pro Phe Leu Glu Leu Lys Ile Arg Ala Asn Glu Pro Gly Ala	
	215 220 225 230	
5	GGC CGG GCC AGG AGG AGG ACC CCC ACC TGT GAG CCT GCG ACC CCC TTA	955
	Gly Arg Ala Arg Arg Arg Thr Pro Thr Cys Glu Pro Ala Thr Pro Leu	
	235 240 245	
	TGT TGC AGG CGA GAC CAT TAC GTA GAC TTC CAG GAA CTG GGA TGG CGG	1003
	Cys Cys Arg Arg Asp His Tyr Val Asp Phe Gln Glu Leu Gly Trp Arg	
	250 255 260	
10	GAC TGG ATA CTG CAG CCC GAG GGG TAC CAG CTG AAT TAC TGC AGT GGG	1051
	Asp Trp Ile Leu Gln Pro Glu Gly Tyr Gln Leu Asn Tyr Cys Ser Gly	
	265 270 275	
	CAG TGC CCT CCC CAC CTG GCT GGC AGC CCA GGC ATT GCT GCC TCT TTC	1099
	Gln Cys Pro Pro His Leu Ala Gly Ser Pro Gly Ile Ala Ala Ser Phe	
15	280 285 290	
	CAT TCT GCC GTC TTC AGC CTC CTC AAA GCC AAC AAT CCT TGG CCT GCC	1147
	His Ser Ala Val Phe Ser Leu Leu Lys Ala Asn Asn Pro Trp Pro Ala	
	295 300 305 310	
	AGT ACC TCC TGT TGT GTC CCT ACT GCC CGA AGG CCC CTC TCT CTC CTC	1195
20	Ser Thr Ser Cys Cys Val Pro Thr Ala Arg Arg Pro Leu Ser Leu Leu	
	315 320 325	
	TAC CTG GAT CAT AAT GGC AAT GTG GTC AAG ACG GAT GTG CCA GAT ATG	1243
	Tyr Leu Asp His Asn Gly Asn Val Val Lys Thr Asp Val Pro Asp Met	
	330 335 340	
25	GTG GTG GAG GCC TGT GGC TGC AGC TAGCAAGAGG ACCTGGGGCT TTGGAGTGAA G	1298
	Val Val Glu Ala Cys Gly Cys Ser	
	345 350	
	AGACCAAGAT GAAGTTTCCC AGGCACAGGG CATCTGTGAC TGGAGGCATC AGATTCCTGA	1358
	TCCACACCCC AACCCAACAA CCACCTGGCA ATATGACTCA CTTGACCCCT ATGGGACCCA	1418
30	AATGGGCACT TTCTTGCTCG AGACTCTGGC TTATTCCAGG TTGGCTGATG TGTTGGGAGA	1478
	TGGGTAAAGC GTTTCTTCTA AAGGGGTCTA CCCAGAAAGC ATGATTTCCT GCCCTAAGTC	1538
	CTGTGAGAAG ATGTCAGGGA CTAGGGAGGG AGGGAGGGAA GGCAGAGAAA AATTACTTAG	1598
	CCTCTCCCAA GATGAGAAAG TCCTCAAGTG AGGGGAGGAG GAAGCAGATA GATGGTCCAG	1658

-43-

CAGGCTTGAA GCAGGGTAAG CAGGCTGGCC CAGGGTAAGG GCTGTTGAGG TACCTTAAGG 1718  
 GAAGGTCAAG AGGGAGATGG GCAAGGCGCT GAGGGAGGAT GCTTAGGGGA CCCCCAGAAA 1778  
 CAGGAGTCAG GAAAATGAGG CACTAAGCCT AAGAAGTTCC CTGGTTTTTC CCAGGGGACA 1838  
 GGACCCACTG GGAGACAAGC ATTTATACTT TCTTTCTTCT TTTTATTTT TTTGAGATCG 1898  
 5 AGTCTCGCTC TGTCACCAGG CTGGAGTGCA GTGACACGAT CTTGGCTCAC TGCAACCTCC 1958  
 GTCTCCTGGG TTCAAGTGAT TCTTCTGCCT CAGCCTCCCG AGCAGCTGGG ATTACAGGCG 2018  
 CCCACTAATT TTTGTATTCT TAGTAGAAAC GAGGTTTCAA CATGTTGGCC AGGATGGTCT 2078  
 CAATCTCTTG ACCTCTTGAT CCACCCGACT TGGCCTCCCG AAGTGATGAG ATTATAGGCG 2138  
 TGAGCCACCG CGCCTGGCTT ATACTTTCTT AATAAAAAGG AGAAAGAAAA TCAACAAATG 2198  
 10 TGAGTCATAA AGAAGGGTTA GGGTGATGGT CCAGAGCAAC AGTTCTTCAA GTGTACTCTG 2258  
 TAGGCTTCTG GGAGGTCCCT TTTCAGGGGT GTCCACAAAG TCAAAGCTAT TTTCATAATA 2318  
 ATACTAACAT GTTATTTGCC TTTTGAATTC TCATTATCTT AAAATTGTAT TGTGGAGTTT 2378  
 TCCAGAGGCC GTGTGACATG TGATTACATC ATCTTCTGA C 2419

(2) INFORMATION FOR SEQ ID NO:14:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 350 amino acids
- (B) TYPE: amino acids
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(x1) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Met Arg Leu Pro Asp Val Gln Leu Trp Leu Val Leu Leu Trp Ala Leu  
 1 5 10 15

Val Arg Ala Gln Gly Thr Gly Ser Val Cys Pro Ser Cys Gly Gly Ser  
 20 25 30

-44-

5

Lys Leu Ala Pro Gln Ala Glu Arg Ala Leu Val Leu Glu Leu Ala Lys  
35 40 45

Gln Gln Ile Leu Asp Gly Leu His Leu Thr Ser Arg Pro Arg Ile Thr  
50 55 60

His Pro Pro Pro Gln Ala Ala Leu Thr Arg Ala Leu Arg Arg Leu Gln  
65 70 75 80

Pro Gly Ser Val Ala Pro Gly Asn Gly Glu Glu Val Ile Ser Phe Ala  
85 90 95

10

Thr Val Thr Asp Ser Thr Ser Ala Tyr Ser Ser Leu Leu Thr Phe His  
100 105 110

Leu Ser Thr Pro Arg Ser His His Leu Tyr His Ala Arg Leu Trp Leu  
115 120 125

His Val Leu Pro Thr Leu Pro Gly Thr Leu Cys Leu Arg Ile Phe Arg  
130 135 140

15

Trp Gly Pro Arg Arg Arg Arg Gln Gly Ser Arg Thr Leu Leu Ala Glu  
145 150 155 160

His His Ile Thr Asn Leu Gly Trp His Thr Leu Thr Leu Pro Ser Ser  
165 170 175

20

Gly Leu Arg Gly Glu Lys Ser Gly Val Leu Lys Leu Gln Leu Asp Cys  
180 185 190

Arg Pro Leu Glu Gly Asn Ser Thr Val Thr Gly Gln Pro Arg Arg Leu  
195 200 205

Leu Asp Thr Ala Gly His Gln Gln Pro Phe Leu Glu Leu Lys Ile Arg  
210 215 220

25

Ala Asn Glu Pro Gly Ala Gly Arg Ala Arg Arg Arg Thr Pro Thr Cys  
225 230 235 240

Glu Pro Ala Thr Pro Leu Cys Cys Arg Arg Asp His Tyr Val Asp Phe  
245 250 255

30

Gln Glu Leu Gly Trp Arg Asp Trp Ile Leu Gln Pro Glu Gly Tyr Gln  
260 265 270

Leu Asn Tyr Cys Ser Gly Gln Cys Pro Pro His Leu Ala Gly Ser Pro  
275 280 285

5  
10  
15  
20  
25  
30

-45-

Gly Ile Ala Ala Ser Phe His Ser Ala Val Phe Ser Leu Leu Lys Ala  
290 295 300

Asn Asn Pro Trp Pro Ala Ser Thr Ser Cys Cys Val Pro Thr Ala Arg  
305 310 315 320

5 Arg Pro Leu Ser Leu Leu Tyr Leu Asp His Asn Gly Asn Val Val Lys  
325 330 335

Thr Asp Val Pro Asp Met Val Val Glu Ala Cys Gly Cys Ser  
340 345 350

55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

**CLAIMS**

1. Substantially pure growth differentiation factor-12 (GDF-12) and functional fragments thereof.
2. An isolated polynucleotide sequence encoding the GDF-12 polypeptide of claim 1.
3. The polynucleotide of claim 2, wherein the GDF-12 is selected from the group consisting of:
  - a. SEQ ID NO:13, wherein T can also be U;
  - b. nucleic acid sequences complementary to SEQ ID NO:13; and
  - c. fragments of a. or b. that are at least 15 bases in length and that will selectively hybridize to DNA which encodes the GDF-12 protein of SEQ ID NO:14; and
4. The polynucleotide of claim 2, wherein the polynucleotide is isolated from a mammalian cell.
5. The polynucleotide of claim 4, wherein the mammalian cell is selected from the group consisting of mouse, rat, and human cell.
6. An expression vector including the polynucleotide of claim 2.
7. The vector of claim 6, wherein the vector is a plasmid.
8. The vector of claim 6, wherein the vector is a virus.
9. A host cell stably transformed with the vector of claim 6.
10. The host cell of claim 9, wherein the cell is prokaryotic.

-47-

11. The host cell of claim 9, wherein the cell is eukaryotic.
12. Antibodies that bind to the polypeptide of claim 1 or fragments thereof.
13. The antibodies of claim 12, wherein the antibodies are polyclonal.
14. The antibodies of claim 12, wherein the antibodies are monoclonal.
15. A method of detecting a cell proliferative disorder comprising contacting the antibody of claim 12 with a specimen of a subject suspected of having a GDF-12 associated disorder and detecting binding of the antibody.
16. The method of claim 15, wherein the cell is a liver cell.
17. The method of claim 15, wherein the detecting is *in vivo*.
18. The method of claim 17, wherein the antibody is detectably labeled.
19. The method of claim 18, wherein the detectable label is selected from the group consisting of a radioisotope, a fluorescent compound, a bioluminescent compound and a chemiluminescent compound.
20. The method of claim 15, wherein the detection is *in vitro*.
21. The method of claim 20, wherein the antibody is detectably labeled.
22. The method of claim 21, wherein the label is selected from the group consisting of a radioisotope, a fluorescent compound, a bioluminescent compound, a chemoluminescent compound and an enzyme

-48-

23. A method of treating a cell proliferative disorder associated with expression of GDF-12, comprising contacting the cells with a reagent which suppresses the GDF-12 activity.
24. The method of claim 23, wherein the reagent is an anti-GDF-12 antibody.
25. The method of claim 23, wherein the reagent is a GDF-12 antisense sequence.
26. The method of claim 23, wherein the cell is a liver cell.
27. The method of claim 23, wherein the reagent which suppresses GDF-12 activity is introduced to a cell using a vector.
28. The method of claim 27, wherein the vector is a colloidal dispersion system.
29. The method of claim 28, wherein the colloidal dispersion system is a liposome.
30. The method of claim 29, wherein the liposome is essentially target specific.
31. The method of claim 30, wherein the liposome is anatomically targeted.
32. The method of claim 31, wherein the liposome is mechanistically targeted.
33. The method of claim 32, wherein the mechanistic targeting is passive.
34. The method of claim 32, wherein the mechanistic targeting is active.

35. The method of claim 34, wherein the liposome is actively targeted by coupling with a moiety selected from the group consisting of a sugar, a glycolipid, and a protein.
36. The method of claim 35, wherein the protein moiety is an antibody.
37. The method of claim 36, wherein the vector is a virus.
38. The method of claim 37, wherein the virus is an RNA virus.
39. The method of claim 38, wherein the RNA virus is a retrovirus.
40. The method of claim 39, wherein the retrovirus is essentially target specific.
41. The method of claim 40, wherein a moiety for target specificity is encoded by a polynucleotide inserted into the retroviral genome.
42. The method of claim 40, wherein a moiety for target specificity is selected from the group consisting of a sugar, a glycolipid, and a protein.
43. The method of claim 42, wherein the protein is an antibody.



GROWTH DIFFERENTIATION FACTOR-12

ABSTRACT OF THE DISCLOSURE

Growth differentiation factor-12 (GDF-12) is disclosed along with its polynucleotide sequence and  
5 amino acid sequence. Also disclosed are diagnostic and therapeutic methods of using the GDF-12 polypeptide and polynucleotide sequences.

221943.B11

221943.B11



ovary  
liver  
muscle  
testis  
spleen  
intestine  
pancreas  
seminal vesicle  
kidney  
brain  
thymus  
lung  
heart

FIG. 1

09301653-073799

```

1  CGGGCCAGGAGGAGGACCCCCACCTGTGAGCCTGCGACCCCTTATGTTGCAGGCGAGAC 60
   R A R R R T P T C E P A T P L C C R R D
61  CATTACGTAGACTTCCAGGAAGTGGGATGGCGGACTGGATACTGCAGCCCGAGGGGTAC 120
   H Y V D F Q E L G W R D W I L Q P E G Y
121 CAGCTGAATTACTGCAGTGGGCAGTGGCCTCCCCACCTGGCTGGCAGCCCAGGCATTGCT 180
   Q L N Y C S G Q C P P H L A G S P G I A
181 GCCTCTTTCATTCTGCCGTCTTCAGCCTCCTCAAAGCCAACAATCCTTGGCCTGCCAGT 240
   A S F H S A V F S L L K A N N P W P A S
241 ACCTCCTGTTGTGTCCCTACTGCCCCAAGGCCCTCTCTCTCTACCTGGATCATAAT 300
   T S C C V P T A R R P L S L L Y L D H N
301 GGCAATGTGGTCAAGACGGATGTGCCAGATATGGTGGTGGAGGCCTGTGGCTGCAGCTAG 360
   G N V V K T D V P D M V V E A C G C S *

```

FIG. 2

<u>Family member</u>	<u>% identity with GDF-12</u>
GDF-1	43
GDF-3	36
GDF-5	36
GDF-6	39
GDF-7	42
GDF-9	30
BMP-3	37
BMP-2	43
BMP-4	42
Vgr-1	41
OP-1	40
BMP-5	38
OP-2	39
MIS	30
Inhibin- $\alpha$	27
Inhibin- $\beta$ A	47
Inhibin- $\beta$ B	50
Nodal	38
GDNF	21
TGF- $\beta$ 1	36
TGF- $\beta$ 2	36
TGF- $\beta$ 3	41

FIG. 4

1	GAGCTGTGAGGGTCAAGCACAGCTATCCATCAGATGATCTACTTTTCAGCCTTCCTGAGTC	60	3/3
61	CCAGACAATAGAAGACAGGTGGCTGTACCCCTTGGCCAAGGGTAGGTGTGGCAGTGGTGTCT	120	
121	TGCTGTCACTGTGCCCTCATTTGGCCCCCAGCAATCAGACTCAACAGACGGAGCAACTGCC	180	
181	ATCCGAGGCTCCTGAACCAGGGCCATTACCAGGAGCATGCGGCTCCCTGATGTCCAGCT	240	
	M R L P D V Q L		
241	CTGGCTGGTGCTGCTGTGGGCACTGGTGGCAGCACAGGGGACAGGGTCTGTGTGTCCCTC	300	
	W L V L L W A L V R A Q G T G S V C P S		
301	CTGTGGGGGCTCCAAACTGGCACCCCAAGCAGAACGAGCTCTGGTGCTGGAGCTAGCCAA	360	
	C G G S K L A P Q A E R A L V L E L A K		
361	GCAGCAAATCCTGGATGGGTGACCTGACCAGTCGTCCCAGAATAACTCATCCTCCACC	420	
	Q Q I L D G L H L T S R P R I T H P P P		
421	CCAGGCAGCGCTGACCAGAGCCCTCCGGAGACTACAGCCAGGGAGTGTGGCTCCAGGGAA	480	
	Q A A L T R A L R R L Q P G S V A P G N		
481	TGGGGAGGAGGTTCATCAGCTTTGCTACTGTACAGACTCCACTTCAGCCTACAGCTCCCT	540	
	G E E V I S F A T V T D S T S A Y S S L		
541	GCTCACTTTTACCTGTCCACTCCTCGGTCCCACCACCTGTACCATGCCCGCCTGTGGCT	600	
	L T F H L S T P R S H H L Y H A R L W L		
601	GCACGTGCTCCCCACCCTTCTGGCACTCTTGTCTGAGGATCTTCCGATGGGGACCAAG	660	
	H V L P T L P G T L C L R I F R W G P R		
661	GAGGAGCGCCAAGGGTCCCGCACTCTCCTGGCTGAGCACCACATCACCAACCTGGGCTG	720	
	R R R Q G S R T L L A E H H I T N L G W		
721	GCATACCTTAACCTCTGCCCTCTAGTGGCTTGAGGGGTGAGAAGTCTGGTGTCTGAAACT	780	
	H T L T L P S S G L R G E K S G V L K L		
781	GCAACTAGACTGCAGACCCCTAGAAGGCAACAGCACAGTTACTGGACAACCGAGGCGGCT	840	
	Q L D C R P L E G <b>N S T</b> V T G Q P R R L		
841	CTTGACACAGCAGGACACCAGCAGCCCTTCTAGAGCTTAAGATCCGAGCCAATGAGCC	900	
	L D T A G H Q Q P F L E L K I R A N E P		
901	TGGAGCAGGCGGGGCCAGGAGGAGGACCCCACTGTGAGCCTGCGACCCCTTATGTTG	960	
	G A G <b>R A E R R R</b> T P T C E P A T P L C C		
961	CAGGCGAGACCATTACGTAGACTTCCAGGAAGTGGGATGGCGGACTGGATACTGCAGCC	1020	
	R R D H Y V D F Q E L G W R D W I L Q P		
1021	CGAGGGTACCAGCTGAATTACTGCAGTGGGCACTGCCCTCCCCACCTGGCTGGCAGCCC	1080	
	E G Y Q L N Y C S G Q C P P H L A G S P		
1081	AGGCATTGCTGCCTCTTTCCATTCTGCCGTCTTCAGCCTCCTCAAAGCCAACAATCCTTG	1140	
	G I A A S F H S A V F S L L K A N N P W		
1141	GCCTGCCAGTACCTCCTGTTGTGTCCCTACTGCCCGAAGGCCCTCTCTCTCTCTACCT	1200	
	P A S T S C C V P T A R R P L S L L Y L		
1201	GGATCATAATGGCAATGTGGTCAAGACGGATGTGCCAGATATGGTGGTGGAGGCCTGTGG	1260	
	D H N G N V V K T D V P D M V V E A C G		
1261	CTGCAGCTAGCAAGAGGACCTGGGGCTTTGGAGTGAAGAGACCAAGATGAAGTTTCCCAG	1320	
	C S *		
1321	GCACAGGGCATCTGTGACTGGAGGCATCAGATTCTGTATCCACACCCCAACCCAACAACC	1380	
1381	ACCTGGCAATATGACTCACTTGACCCCTATGGGACCCAAATGGGCACTTTCTTGTCTGAG	1440	
1441	ACTCTGGCTTATTCCAGGTGGCTGATGTGTGGGAGATGGGTAAAGCGTTTCTTCTAAA	1500	
1501	GGGGTCTACCCAGAAAGCATGATTTCTGCCCCAAGTCTGTGAGAAGATGTCAGGGACT	1560	
1561	AGGGAGGGAGGGAGGGAAGGCAGAGAAAAATTACTTAGCCTCTCCCAAGATGAGAAAGTC	1620	
1621	CTCAAGTGAGGGGAGGAGGAGCAGATAGATGGTCCAGCAGGCTTGAAGCAGGGTAAGCA	1680	
1681	GGCTGGCCCCAGGGTAAGGGCTGTTGAGGTACCTTAAGGGAAGGTCAAGAGGGAGATGGGC	1740	
1741	AAGGCGCTGAGGGAGGATGCTTAGGGGACCCCAAGAACAGGAGTCAGGAAAATGAGGCA	1800	
1801	CTAAGCCTAAGAAGTTCCCTGGTTTTTCCCAGGGGACAGGACCCACTGGGAGACAAGCAT	1860	
1861	TTATACTTTCTTTCTTTCTTTTATTTTTTTGAGATCGAGTCTCGCTCTGTACCAGGCT	1920	
1921	GGAGTGCAGTGACACGATCTTGGCTCACTGCAACCTCCGTCTCCTGGGTTCAAGTGATTC	1980	
1981	TTCTGCCCTCAGCCTCCCGAGCAGCTGGGATTACAGGCGCCCACTAATTTTTGTATTCTTA	2040	
2041	GTAGAAACGAGGTTTCAACATGTTGGCCAGGATGGTCTCAATCTCTTGACCTCTTGATCC	2100	
2101	ACCCGACTTGGCCTCCCGAAGTGATGAGATTATAGGCGTGAGCCACCGCGCCTGGCTTAT	2160	
2161	ACTTTCTTAATAAAAAGGAGAAAGAAATCAACAAATGTGAGTCATAAAGAAGGGTTAGG	2220	
2221	GTGATGGTCCAGAGCAACAGTCTTCAAGTGTACTCTGTAGGCTTCTGGAGGCTCCCTTT	2280	
2281	TCAGGGGTGTCCACAAAGTCAAAGCTATTTTCATAATAATACTAACATGTTATTTGCCTT	2340	
2341	TGAATTCTCATTATCTTAAATTTGATTGTGGAGTTTCCAGAGGCCGTGTGACATGTG	2400	
2401	ATTACATCATCTTCTGAC 2419		

FIG. 3

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants: Se-Jin Lee *et al.* Art Unit: 1812  
Serial No.: 08/765,662 Examiner: P. Mertz  
Filed: 4/28/97  
Title: GROWTH DIFFERENTIATION FACTOR-12

Assistant Commissioner for Patents  
Washington DC 20231


**TRANSMITTAL OF CORRECTED DECLARATION**

Enclosed is the executed revised Combined Declaration and Power of Attorney previously filed (unsigned) on April 24, 1998, in response to the Office Action dated February 24, 1998.

No additional fees are believed to be due. Please charge any required fee, or make any credits, to Deposit Account No. 06-1050.

Respectfully submitted,

Date: 6/5/98



Lisa A. Haile, Ph.D.


Reg. No. 38,347

Attorney for Applicants

Fish & Richardson P.C.  
4225 Executive Square, Suite 1400  
La Jolla, CA 92037  
Telephone: 619/678-5070  
Facsimile: 619/678-5099  
55521.LJ1

Date of Deposit 6/8/98

I hereby certify under 37 CFR 1.8(a) that this correspondence is being deposited with the United States Postal Service as **first class mail** with sufficient postage on the date indicated above and is addressed to the Commissioner of Patents and Trademarks, Washington, D.C. 20231.



COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled GROWTH DIFFERENTIATION FACTOR-12, the specification of which

☐ is attached hereto.

☐ was filed on \_\_\_\_\_ as Application Serial No. \_\_\_\_\_  
and was amended on \_\_\_\_\_.

☒ was described and claimed in PCT International Application No. PCT/US95/08745 filed on 7/12/95  
and was amended on .

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose all information I know to be material to patentability in accordance with Title 37, Code of Federal Regulations, §1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed:

COUNTRY	APPLICATION NO.	FILING DATE	PRIORITY CLAIMED
_____	_____	_____	<input type="checkbox"/> Yes <input type="checkbox"/> No

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose all information I know to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56(a) which became available between the filing date of the prior application and the national or PCT international filing date of this application:

U.S. SERIAL NO.	FILING DATE	STATUS
<u>08/274,215</u>	<u>7/13/94</u>	<input checked="" type="checkbox"/> Pending <input type="checkbox"/> Issued <input type="checkbox"/> Abandoned
<u>PCT/US95/08745</u>	<u>7/12/95</u>	<input checked="" type="checkbox"/> Pending <input type="checkbox"/> Issued <input type="checkbox"/> Abandoned

I hereby appoint the following attorneys and/or agents to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith: Lisa A. Haile, Reg. No. 38,347, John R. Wetherell, Reg. No. 31,678, June M. Learn, Reg. No. 31,238, John Land, Reg. No. 29,554..


Address all telephone calls to Lisa A. Haile at telephone number 619/678-5070.

Address all correspondence to Lisa A. Haile, Fish & Richardson P.C., 4225 Executive Square, Suite 1400, La Jolla, CA 92037.

## COMBINED DECLARATION AND POWER OF ATTORNEY CONTINUED

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patents issued thereon.

Full Name of Inventor: Se-Jin Lee

Inventor's Signature: 


Date: 5/20/98

Residence Address: 6711 Chokeberry Road, Baltimore, MD 21205

Citizen of: USA

Post Office Address: 6711 Chokeberry Road, Baltimore, MD 21205

Full Name of Inventor: Aurora F. Esqueda

Inventor's Signature: 

Date: 5/20/98

Residence Address: Apartment 1B, 853 West University Parkway, Baltimore MD 21210

Citizen of: USA

Post Office Address: Apartment 1B, 853 West University Parkway, Baltimore MD 21210

09761655 070299

## COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled GROWTH DIFFERENTIATION FACTOR-12, the specification of which

☐ is attached hereto.

☐ was filed on \_\_\_\_\_ as Application Serial No. \_\_\_\_\_  
and was amended on \_\_\_\_\_.

☒ was described and claimed in PCT International Application No. PCT/US95/08745 filed on 7/12/95  
and was amended on .

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose all information I know to be material to patentability in accordance with Title 37, Code of Federal Regulations, §1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed:

COUNTRY	APPLICATION NO.	FILING DATE	PRIORITY CLAIMED
<u>PCT</u>	<u>PCT/US95/08754</u>	<u>7/12/95</u>	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose all information I know to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56(a) which became available between the filing date of the prior application and the national or PCT international filing date of this application:

U.S. SERIAL NO.	FILING DATE	STATUS
<u>08/274,215</u>	<u>7/13/94</u>	<input checked="" type="checkbox"/> Pending <input type="checkbox"/> Issued <input type="checkbox"/> Abandoned
<u>08/311,370</u>	<u>9/26/94</u>	<input checked="" type="checkbox"/> Pending <input type="checkbox"/> Issued <input type="checkbox"/> Abandoned

I hereby appoint the following attorneys and/or agents to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith: Lisa A. Haile, Reg. No. 38,347, John R. Wetherell, Reg. No. 31,678, June M. Learn, Reg. No. 31,238, John Land, Reg. No. 29,554..

Address all telephone calls to Lisa A. Haile at telephone number 619/678-5070.

Address all correspondence to Lisa A. Haile, Fish & Richardson P.C., 4225 Executive Square, Suite 1400, La Jolla, CA 92037.


I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the



## COMBINED DECLARATION AND POWER OF ATTORNEY CONTINUED

knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patents issued thereon.

Full Name of Inventor: Se-Jin Lee

Inventor's Signature: 


Date: 4/23/97

Residence Address: 6711 Chokeberry Road, Baltimore, MD 21205

Citizen of: USA

Post Office Address: 6711 Chokeberry Road, Baltimore, MD 21205

Full Name of Inventor: Aurora F. Esquela

Inventor's Signature: 

Date: 4/23/97

Residence Address: Apartment 1B, 853 West University Parkway, Baltimore MD 21210

Citizen of: USA

Post Office Address: Apartment 1B, 853 West University Parkway, Baltimore MD 21210

09363660